


Anti-Beta Catenin, mouse monoclonal (BS31)

BSH-2014-100 (0.1 ml), BSH-2014-1 (1 ml)


Clonality:	Mouse monoclonal antibody
Clone:	BS31
Application:	IHC-P (1:100 – 1:400), IHC-Fro
Species Reactivity:	Human
Control tissues:	Appendix, liver, tonsil
Buffer:	TRIS with 0.03% sodium azide, pH 7.2
Storage:	Store at 4°C



Liver section has been stained using Beta-Catenin optibody (Clone: BS31) with 1:200 dilution. Hepatocytes and bile ducts have strong membranous label.

Description

Beta-Catenin is a member of catenin family together with alpha and gamma catenin. It mediates cell-cell adhesion with cadherins and it is key regulatory protein in signaling through the WNT pathway. Beta catenin has a role in cellular proliferation, differentiation and development. Mutations in beta catenin gene (CTNNB1) leads accumulation of the beta catenin protein in cytoplasm and nucleus in different type of tumors eg. endometrial carcinoma and desmoid tumors. This antibody is useful in differentiation diagnostic of tumors.




Tonsil section has been stained using Beta-Catenin optibody (Clone: BS31) with 1:200 dilution. Epithelial cells have strong label. Follicular dendritic cells and vascular endothelial cells have label in tonsil.

Protocol

1. Deparaffinize and rehydrate tissue section
2. Wash: aqua dest, 2×5 min
3. Pre-treatment: PT-module HIER pH 9.0 (20min at 98°C)
4. H₂O₂ (concentration 3%), 10 min
5. Wash: PBS or TBS buffer, 2×5 min
6. Primary antibody diluted as recommended, 30 min
7. Wash: PBS or TBS buffer, 2×5 min
8. One step HRP-polymer detection, 30 min
9. Wash: PBS or TBS buffer, 2×5 min
10. DAB Substrate, 8 min
11. Wash: aqua dest, 2×2 min
12. Counterstain, dehydrate and coverslip

Dilution of concentrated antibody depends on the pre-treatment method and detection system used. Above protocol used in Optibodies evaluation and is meant as a reference. Final working dilution and protocol applied needs to be determined by the user always.



Ductal breast carcinoma section has been stained using Beta-Catenin optibody (Clone: BS31) with 1:200 dilution. Ductals carcinoma cells have strong membranous staining pattern, without nuclear accumulation of beta-catenin.