


Anti-CD14, mouse monoclonal (BS9)

BSH-7019-100 (0.1 ml), BSH-7019-1 (1 ml)




Clonality:	Mouse monoclonal antibody
Clone:	BS9
Application:	IHC-P (1:100 – 1:400), IHC-Fro
Species Reactivity:	Human
Control tissues:	Tonsil, liver, appendix
Buffer:	TRIS with 0.03% sodium azide, pH 7.2
Storage:	Store at 4°C



Tonsil section have been stained using CD14 optibody (Clone: BS9) with 1:200 dilution. Follicular dendritic cells have strong label as well as macrophages in perifollicular area.

Description

CD14 antigen is a GPI-linked glycoprotein with a molecular weight of 55kD. It is expressed on cells of the myelomonocytic lineage including monocytes, macrophages and Langerhans cells. Low expression is observed on neutrophils and on human B cells. CD14 antigen is a receptor for bacterial lipopolysaccharide (LPS, endotoxin) and the lipopolysaccharide binding protein (LBP). LBP and CD14 antigen serve two physiological roles. These proteins act as opsonin and opsonic receptor, respectively, to promote the phagocytic uptake of bacteria or LPS coated particles by macrophages.




Tonsil section have been stained using CD14 optibody (Clone: BS9) with 1:200 dilution. Follicular dendritic cells have strong label as well as macrophages in perifollicular area.

Protocol

1. Deparaffinize and rehydrate tissue section
2. Wash: aqua dest, 2x5 min
3. Pre-treatment: PT-module HIER pH 9.0 (20min at 98°C)
4. H₂O₂ (concentration 3%), 10 min
5. Wash: PBS or TBS buffer, 2x5 min
6. Primary antibody diluted as recommended, 30 min
7. Wash: PBS or TBS buffer, 2x5 min
8. One step HRP-polymer detection, 30 min
9. Wash: PBS or TBS buffer, 2x5 min
10. DAB Substrate, 8 min
11. Wash: aqua dest, 2x2 min
12. Counterstain, dehydrate and coverslip

Dilution of concentrated antibody depends on the pre-treatment method and detection system used. Above protocol used in Optibodies evaluation and is meant as a reference. Final working dilution and protocol applied needs to be determined by the user always.



Liver section have been stained using CD14 optibody (Clone: BS9) with 1:200 dilution. Kupffer cells have strong to moderate label.