

Anti-CD10, mouse monoclonal (BS1)

BSH-7021-100 (0,1ml), BSH-7021-1 (1 ml)

Clonality:	Mouse monoclonal antibody
Clone:	BS1
Application:	IHC-P (1:100 – 1:400)
Species Reactivity:	Human
Control tissues:	Tonsil, liver, kidney
Buffer:	TRIS with 0.03% sodium azide, pH 7,2
Storage:	Store at 4°C



Description:

CD10 is a 100kDa glycoprotein, also designated Common Acute Lymphocytic Leukemia Antigen (CALLA). It is a cell surface enzyme with neutral metalloendopeptidase activity which inactivates a variety of biologically active peptides. CD10 is expressed on the cells of lymphoblastic, Burkitt's, and follicular germinal center lymphomas, and on cells from patients with chronic myelocytic leukemia (CML). It is also expressed on the surface of normal early lymphoid progenitor cells, immature B cells within adult bone marrow and germinal center B cells within lymphoid tissue. CD10 is also present on breast myoepithelial cells, bile canaliculi, fibroblasts, with especially high expression on the brush border of kidney and gut epithelial cells. CD10 is useful especially in the classification of B-cell leukemias and lymphomas.


Protocol

After paraffin removing and rehydration:


1. Pre-treatment: PT-module HIER pH9 (20min at 98°C)
2. Wash (TBS-Tween in all washing steps)
3. Primary antibody: CD10 1:100 – 1:400, 30 min.
4. Wash
5. Peroxidase blocking (3% H₂O₂), 10 min.
6. Wash
7. One step HRP-polymer detection, 30 min
8. Wash x2
9. DAB-Substrate, 10 min
10. Aqua
11. CuSO₄ -post enhancement, 5 min
12. Aqua

Counter staining, Bluing, dehydration, clearing, and mounting.


Dilution of concentrated antibody depends on the pre-treatment method and detection system used. Above protocol used in Optibodies evaluation and is meant as a reference. Final working dilution and protocol applied needs to be determined by the user always.



Tonsil section have been stained using CD10 antibody (Clone: BS1) with 1:300 dilution. Follicular B-cells have moderate to strong membranous staining pattern.



Kidney section have been stained using CD10 antibody (Clone: BS1) with 1:300 dilution. Proximal tubules and glomerulus have strong staining reaction.



Liver section have been stained using CD10 antibody (Clone: BS1) with 1:300 dilution. Bile canaliculi have strong label without staining in hepatocytes.